



Pairwise significance tests for the proportions or medians presented in Figures 2-4

For proportions, the significance test in Tables A and C follows Fleiss et al. (2003: 188-189, cited in the article). The test compares several (here 2) proportions, based on a χ^2 -test. For medians in Tables B and D, the non-parametric Mann-Whitney U-test was used. It is described, for example, in Sokal & Rohlf (2012: 446-452, cited in the article). The probability refers to the null hypothesis that the two proportions or two medians come from the same statistical populations. Since there is a relatively large number of simultaneous pairwise comparisons in each table, the threshold level for concluding significant differences (typically 0.05) should be adjusted: Using the Bonferroni method, mentioned in the article, the adjusted significance level in Tables A and B (with 10 proportions or medians) is $0.05/10 = \mathbf{0.005}$. The adjusted significance level in Table C and D (with 4 proportions or medians) is $0.05/4 \approx \mathbf{0.01}$. The sequence of species in the tables is in alphabetical order.

Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement 1.



Table A. Pairwise significance tests between survival proportions (Figure 2 top)

First species	Second species	Probability
<i>Brosimum alicastrum</i>	<i>Calophyllum brasiliense</i>	0.136
<i>Brosimum alicastrum</i>	<i>Ceiba pentandra</i>	0.000
<i>Brosimum alicastrum</i>	<i>Colubrina arborescens</i>	0.000
<i>Brosimum alicastrum</i>	<i>Diphysa americana</i>	0.000
<i>Brosimum alicastrum</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Brosimum alicastrum</i>	<i>Handroanthus chrysanthus</i>	0.000
<i>Brosimum alicastrum</i>	<i>Pimenta dioica</i>	0.000
<i>Brosimum alicastrum</i>	<i>Pouteria sapota</i>	0.385
<i>Brosimum alicastrum</i>	<i>Tabebuia rosea</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Ceiba pentandra</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Colubrina arborescens</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Diphysa americana</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Handroanthus chrysanthus</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Pimenta dioica</i>	0.014
<i>Calophyllum brasiliense</i>	<i>Pouteria sapota</i>	0.530
<i>Calophyllum brasiliense</i>	<i>Tabebuia rosea</i>	0.000
<i>Ceiba pentandra</i>	<i>Colubrina arborescens</i>	0.009
<i>Ceiba pentandra</i>	<i>Diphysa americana</i>	0.001
<i>Ceiba pentandra</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Ceiba pentandra</i>	<i>Handroanthus chrysanthus</i>	0.017
<i>Ceiba pentandra</i>	<i>Pimenta dioica</i>	0.000
<i>Ceiba pentandra</i>	<i>Pouteria sapota</i>	0.000
<i>Ceiba pentandra</i>	<i>Tabebuia rosea</i>	0.000
<i>Colubrina arborescens</i>	<i>Diphysa americana</i>	0.471
<i>Colubrina arborescens</i>	<i>Enterolobium cyclocarpum</i>	0.140
<i>Colubrina arborescens</i>	<i>Handroanthus chrysanthus</i>	0.825
<i>Colubrina arborescens</i>	<i>Pimenta dioica</i>	0.000
<i>Colubrina arborescens</i>	<i>Pouteria sapota</i>	0.000
<i>Colubrina arborescens</i>	<i>Tabebuia rosea</i>	0.226
<i>Diphysa americana</i>	<i>Enterolobium cyclocarpum</i>	0.450
<i>Diphysa americana</i>	<i>Handroanthus chrysanthus</i>	0.347
<i>Diphysa americana</i>	<i>Pimenta dioica</i>	0.004
<i>Diphysa americana</i>	<i>Pouteria sapota</i>	0.000
<i>Diphysa americana</i>	<i>Tabebuia rosea</i>	0.625
<i>Enterolobium cyclocarpum</i>	<i>Handroanthus chrysanthus</i>	0.090
<i>Enterolobium cyclocarpum</i>	<i>Pimenta dioica</i>	0.036
<i>Enterolobium cyclocarpum</i>	<i>Pouteria sapota</i>	0.000
<i>Enterolobium cyclocarpum</i>	<i>Tabebuia rosea</i>	0.789
<i>Handroanthus chrysanthus</i>	<i>Pimenta dioica</i>	0.000
<i>Handroanthus chrysanthus</i>	<i>Pouteria sapota</i>	0.000
<i>Handroanthus chrysanthus</i>	<i>Tabebuia rosea</i>	0.155
<i>Pimenta dioica</i>	<i>Pouteria sapota</i>	0.002
<i>Pimenta dioica</i>	<i>Tabebuia rosea</i>	0.018
<i>Pouteria sapota</i>	<i>Tabebuia rosea</i>	0.000

Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement I.



Table B. Pairwise significance tests between medians of height increment (Figure 2 bottom)

First species	Second species	Probability
<i>Brosimum alicastrum</i>	<i>Calophyllum brasiliense</i>	0.002
<i>Brosimum alicastrum</i>	<i>Ceiba pentandra</i>	0.000
<i>Brosimum alicastrum</i>	<i>Colubrina arborescens</i>	0.000
<i>Brosimum alicastrum</i>	<i>Diphysa americana</i>	0.029
<i>Brosimum alicastrum</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Brosimum alicastrum</i>	<i>Handroanthus chrysanthus</i>	0.283
<i>Brosimum alicastrum</i>	<i>Pimenta dioica</i>	0.669
<i>Brosimum alicastrum</i>	<i>Pouteria sapota</i>	0.159
<i>Brosimum alicastrum</i>	<i>Tabebuia rosea</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Ceiba pentandra</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Colubrina arborescens</i>	0.026
<i>Calophyllum brasiliense</i>	<i>Diphysa americana</i>	0.404
<i>Calophyllum brasiliense</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Handroanthus chrysanthus</i>	0.094
<i>Calophyllum brasiliense</i>	<i>Pimenta dioica</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Pouteria sapota</i>	0.000
<i>Calophyllum brasiliense</i>	<i>Tabebuia rosea</i>	0.000
<i>Ceiba pentandra</i>	<i>Colubrina arborescens</i>	0.000
<i>Ceiba pentandra</i>	<i>Diphysa americana</i>	0.000
<i>Ceiba pentandra</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Ceiba pentandra</i>	<i>Handroanthus chrysanthus</i>	0.000
<i>Ceiba pentandra</i>	<i>Pimenta dioica</i>	0.000
<i>Ceiba pentandra</i>	<i>Pouteria sapota</i>	0.000
<i>Ceiba pentandra</i>	<i>Tabebuia rosea</i>	0.000
<i>Colubrina arborescens</i>	<i>Diphysa americana</i>	0.001
<i>Colubrina arborescens</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Colubrina arborescens</i>	<i>Handroanthus chrysanthus</i>	0.000
<i>Colubrina arborescens</i>	<i>Pimenta dioica</i>	0.000
<i>Colubrina arborescens</i>	<i>Pouteria sapota</i>	0.000
<i>Colubrina arborescens</i>	<i>Tabebuia rosea</i>	0.070
<i>Diphysa americana</i>	<i>Enterolobium cyclocarpum</i>	0.000
<i>Diphysa americana</i>	<i>Handroanthus chrysanthus</i>	0.292
<i>Diphysa americana</i>	<i>Pimenta dioica</i>	0.000
<i>Diphysa americana</i>	<i>Pouteria sapota</i>	0.000
<i>Diphysa americana</i>	<i>Tabebuia rosea</i>	0.000
<i>Enterolobium cyclocarpum</i>	<i>Handroanthus chrysanthus</i>	0.000
<i>Enterolobium cyclocarpum</i>	<i>Pimenta dioica</i>	0.000
<i>Enterolobium cyclocarpum</i>	<i>Pouteria sapota</i>	0.000
<i>Enterolobium cyclocarpum</i>	<i>Tabebuia rosea</i>	0.000
<i>Handroanthus chrysanthus</i>	<i>Pimenta dioica</i>	0.030
<i>Handroanthus chrysanthus</i>	<i>Pouteria sapota</i>	0.019
<i>Handroanthus chrysanthus</i>	<i>Tabebuia rosea</i>	0.000
<i>Pimenta dioica</i>	<i>Pouteria sapota</i>	0.215
<i>Pimenta dioica</i>	<i>Tabebuia rosea</i>	0.000
<i>Pouteria sapota</i>	<i>Tabebuia rosea</i>	0.000

Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement 1.



Table C. Pairwise significance tests between survival proportions (Figures 3 and 4 left)

Species	First dose (g)	Second dose (g)	Probability
<i>Brosimum alicastrum</i>	0	150	0.443
<i>Brosimum alicastrum</i>	0	375	0.474
<i>Brosimum alicastrum</i>	0	750	0.810
<i>Brosimum alicastrum</i>	150	375	0.145
<i>Brosimum alicastrum</i>	150	750	0.316
<i>Brosimum alicastrum</i>	375	750	0.628
<i>Calophyllum brasiliense</i>	0	150	0.046
<i>Calophyllum brasiliense</i>	0	375	0.002
<i>Calophyllum brasiliense</i>	0	750	0.000
<i>Calophyllum brasiliense</i>	150	375	0.220
<i>Calophyllum brasiliense</i>	150	750	0.088
<i>Calophyllum brasiliense</i>	375	750	0.594
<i>Ceiba pentandra</i>	0	150	0.280
<i>Ceiba pentandra</i>	0	375	0.280
<i>Ceiba pentandra</i>	0	750	0.110
<i>Ceiba pentandra</i>	150	375	1.000
<i>Ceiba pentandra</i>	150	750	0.596
<i>Ceiba pentandra</i>	375	750	0.596
<i>Colubrina arborescens</i>	0	150	0.516
<i>Colubrina arborescens</i>	0	375	0.218
<i>Colubrina arborescens</i>	0	750	0.019
<i>Colubrina arborescens</i>	150	375	0.059
<i>Colubrina arborescens</i>	150	750	0.003
<i>Colubrina arborescens</i>	375	750	0.258
<i>Diphysa americana</i>	0	150	0.003
<i>Diphysa americana</i>	0	375	0.070
<i>Diphysa americana</i>	0	750	0.002
<i>Diphysa americana</i>	150	375	0.243
<i>Diphysa americana</i>	150	750	0.880
<i>Diphysa americana</i>	375	750	0.188
<i>Enterolobium cyclocarpum</i>	0	150	0.776
<i>Enterolobium cyclocarpum</i>	0	375	0.549
<i>Enterolobium cyclocarpum</i>	0	750	0.160
<i>Enterolobium cyclocarpum</i>	150	375	0.378
<i>Enterolobium cyclocarpum</i>	150	750	0.092
<i>Enterolobium cyclocarpum</i>	375	750	0.423
<i>Handroanthus chrysanthus</i>	0	150	0.537
<i>Handroanthus chrysanthus</i>	0	375	0.772
<i>Handroanthus chrysanthus</i>	0	750	0.165
<i>Handroanthus chrysanthus</i>	150	375	0.371
<i>Handroanthus chrysanthus</i>	150	750	0.049
<i>Handroanthus chrysanthus</i>	375	750	0.271
<i>Pimenta dioica</i>	0	150	0.427
<i>Pimenta dioica</i>	0	375	0.006
<i>Pimenta dioica</i>	0	750	0.000
<i>Pimenta dioica</i>	150	375	0.050
<i>Pimenta dioica</i>	150	750	0.002
<i>Pimenta dioica</i>	375	750	0.249
<i>Pouteria sapota</i>	0	150	0.057
<i>Pouteria sapota</i>	0	375	0.008
<i>Pouteria sapota</i>	0	750	0.001
<i>Pouteria sapota</i>	150	375	0.410
<i>Pouteria sapota</i>	150	750	0.104
<i>Pouteria sapota</i>	375	750	0.390
<i>Tabebuia rosea</i>	0	150	0.598
<i>Tabebuia rosea</i>	0	375	0.410
<i>Tabebuia rosea</i>	0	750	0.840
<i>Tabebuia rosea</i>	150	375	0.771
<i>Tabebuia rosea</i>	150	750	0.475
<i>Tabebuia rosea</i>	375	750	0.316

Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement I.



Table D. Pairwise significance tests between medians of height increment (Figures 3 and 4 left)

Species	First dose (g)	Second dose (g)	Probability
<i>Brosimum alicastrum</i>	0	150	0.427
<i>Brosimum alicastrum</i>	0	375	0.925
<i>Brosimum alicastrum</i>	0	750	0.104
<i>Brosimum alicastrum</i>	150	375	0.588
<i>Brosimum alicastrum</i>	150	750	0.422
<i>Brosimum alicastrum</i>	375	750	0.459
<i>Calophyllum brasiliense</i>	0	150	0.034
<i>Calophyllum brasiliense</i>	0	375	0.316
<i>Calophyllum brasiliense</i>	0	750	0.335
<i>Calophyllum brasiliense</i>	150	375	0.390
<i>Calophyllum brasiliense</i>	150	750	0.640
<i>Calophyllum brasiliense</i>	375	750	1.000
<i>Ceiba pentandra</i>	0	150	0.003
<i>Ceiba pentandra</i>	0	375	0.000
<i>Ceiba pentandra</i>	0	750	0.000
<i>Ceiba pentandra</i>	150	375	0.059
<i>Ceiba pentandra</i>	150	750	0.088
<i>Ceiba pentandra</i>	375	750	0.743
<i>Colubrina arborescens</i>	0	150	0.530
<i>Colubrina arborescens</i>	0	375	0.058
<i>Colubrina arborescens</i>	0	750	0.740
<i>Colubrina arborescens</i>	150	375	0.192
<i>Colubrina arborescens</i>	150	750	0.380
<i>Colubrina arborescens</i>	375	750	0.045
<i>Diphysa americana</i>	0	150	0.783
<i>Diphysa americana</i>	0	375	0.004
<i>Diphysa americana</i>	0	750	0.836
<i>Diphysa americana</i>	150	375	0.091
<i>Diphysa americana</i>	150	750	0.915
<i>Diphysa americana</i>	375	750	0.010
<i>Enterolobium cyclocarpum</i>	0	150	0.317
<i>Enterolobium cyclocarpum</i>	0	375	0.538
<i>Enterolobium cyclocarpum</i>	0	750	0.034
<i>Enterolobium cyclocarpum</i>	150	375	0.868
<i>Enterolobium cyclocarpum</i>	150	750	0.106
<i>Enterolobium cyclocarpum</i>	375	750	0.348
<i>Handroanthus chrysanthus</i>	0	150	0.453
<i>Handroanthus chrysanthus</i>	0	375	0.212
<i>Handroanthus chrysanthus</i>	0	750	0.023
<i>Handroanthus chrysanthus</i>	150	375	0.563
<i>Handroanthus chrysanthus</i>	150	750	0.064
<i>Handroanthus chrysanthus</i>	375	750	0.223
<i>Pimenta dioica</i>	0	150	0.076
<i>Pimenta dioica</i>	0	375	0.439
<i>Pimenta dioica</i>	0	750	0.653
<i>Pimenta dioica</i>	150	375	0.451
<i>Pimenta dioica</i>	150	750	0.596
<i>Pimenta dioica</i>	375	750	0.755
<i>Pouteria sapota</i>	0	150	0.001
<i>Pouteria sapota</i>	0	375	0.542
<i>Pouteria sapota</i>	0	750	(n ₂ = 1)
<i>Pouteria sapota</i>	150	375	0.017
<i>Pouteria sapota</i>	150	750	(n ₂ = 1)
<i>Pouteria sapota</i>	375	750	(n ₂ = 1)
<i>Tabebuia rosea</i>	0	150	0.001
<i>Tabebuia rosea</i>	0	375	0.001
<i>Tabebuia rosea</i>	0	750	0.000
<i>Tabebuia rosea</i>	150	375	0.684
<i>Tabebuia rosea</i>	150	750	0.637
<i>Tabebuia rosea</i>	375	750	0.514

Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement 1.



Supplementary material of the original: Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico.

Received: 24 September 2020 Accepted: 15 November 2021 Published: 17 June 2022

This document must be cited as:

Martínez-Bravo, R. D., Ricker, M., Nava-Cruz, Y., & Siebe, C. (2022). Mineral fertilization in tropical reforestation: seedling growth and survival of ten native tree species in Tabasco, Mexico. *Madera y Bosques*, 28(1), e2812269. <https://doi.org/10.21829/myb.2022.2812269>. Supplement 1.